

Nutrition and Food Technology: Open Access

Review Article

Volume: 2.2

Open Access

Antioxidants: Balancing the Good, the Bad and the Ugly

Muhammed Majeed^{1,2}, Kalyanam Nagabhushanam² and Ambar K Choudhury^{1*}¹Sami Labs Limited, Bangalore, India²Sabinsa Corporation-East Windsor, NJ, USA***Corresponding author:** Ambar K Choudhury, Sami Labs Limited, Bangalore-560058, India, Tel: +91-80-28397973; Fax: +91-80-28373035; **E-mail:** ambarchoudhury@samilabs.com**Received date:** 02 Mar 2016; **Accepted date:** 28 Apr 2016; **Published date:** 03 May 2016.**Citation:** Majeed M, Nagabhushanam K, Choudhury AK (2016) Antioxidants: Balancing the Good, the Bad and the Ugly. *Nutr Food Technol* 2(2): doi <http://dx.doi.org/10.16966/2470-6086.123>**Copyright:** © 2016 Majeed M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Antioxidants refer to the molecules that react with oxy radicals not only to delay their function, but also to neutralize them and subsequently reduce oxidative stress and protect us from diseases state. Despite our own endogenous response, several factors contribute to diminish our self-defense mechanism resulting in loss of control in preventing ourselves from free radical damage. Antioxidant supplementation is becoming a high-priority to protect our health and from progression of several life-threatening diseases.

Keywords: Antioxidants; Free radicals; Healthy Food; Phytonutrients/phytochemicals; Supplement**Abbreviations:** ROS: Reactive Oxygen Species; RNS: Reactive Nitrogen Species; DNA: Deoxyribonucleic Acid; MsrA: Methionine Sulfoxide Reductase A; HO: Heme Oxygenase; 4-HNE: 4-Hydroxynonenal; TCA: Tricarboxylic acid; Nrf2: Nuclear Factor Erythroid 2-Related Factor; COX-2: Cyclooxygenase-2; SOD: Superoxide Dismutase; DPPH: 1,1-Diphenyl-2-picryl hydrazyl.

Introduction

Free radicals and antioxidants

Nevertheless, oxygen is profoundly essential for the survival; however, reduction of molecular oxygen sequentially produces oxy radicals such as hydroxyl radicals, hydrogen peroxide, superoxide anion radicals often called as free radicals, reactive oxygen species (ROS) and reactive nitrogen species (RNS) like nitric oxide radical. Though some amount of these free radicals essentially involves keeping us healthy, however, excess of unsuppressed ROS and RNS could be deleterious to our health. Even though our body has its own self-defense mechanism to neutralize oxy radicals through the gainful employment of endogenous anti-oxidants, however, oxidative stress, environmental effect, food-habits as well as aging process, reduce human's auto-immune empowerment. Once these free radicals overpower human's innate anti-oxidant capacity ('state of oxidative stress'), they attack healthy cells and other organs and ultimately damage healthy cellular DNA causing the progression of several life-threatening diseases like cancer [1,2], diabetes mellitus [3,4], rheumatoid arthritis [5], ulcers [6], stroke [7], cardiovascular diseases [8]. Uncontrolled free radicals are responsible for other health disorders such as Parkinson's disease [9], eye diseases [10] inclusive of cataracts [11], noise induced hearing loss [12,13], aging related disorder [14] and disorient humans being from leading healthy life-style.

Antioxidants [15] refer to the molecules that react with oxy and other radicals not only to delay their function, but also to neutralize them and subsequently protect us from oxidative stress [16] and keep us away from other diseases. The major classes of antioxidants are i) enzymatic antioxidants, ii) non-enzymatic antioxidants and iii) metal binding proteins [17] that play the decisive role for controlling our body over oxy-radical and damage from other radicals.

Many methods are now well recognized and are in use for the measurement of antioxidants capacity. Very recently, three consecutive

reports [18-20] elaborated in detail with mechanistic approach on the measurement of antioxidants capacity or its activity.

Enzymatic antioxidants are mainly endogenous as they exist in our body and are very powerful antioxidants. Despite number of enzymes in our body belonging to this group, three super active enzymes are superoxide dismutase, catalase and glutathione. They catalyze the decomposition of superoxide, hydrogen peroxide and organic peroxides and detoxify oxy radicals and convert them finally into oxygen, water and alcohols. Metal binding proteins [17] such as albumin, transferrin, ferritin, lactoferrin are present in our body and protect us from severe damage by trapping free radicals.

Non-enzymatic antioxidants are mostly exogenous as our body hardly produce them and need to be outsourced. They are i) nutritional antioxidants such as vitamin C, vitamin E, carotenoids, ii) mineral constituents like selenium and iii) natural antioxidants: derived from natural sources often known as phytonutrients/phytochemicals.

Antioxidants and Human Health Benefits

Antioxidants in food [21,22]

Dietary foods are substantial source of antioxidants. Antioxidants, present in food are both nutritional like vitamin C, vitamin E, carotenoids and non-nutrients such as polyphenols and others elements. Vitamin C is a water soluble antioxidant and is highly active to neutralize oxy radicals in aqueous phase whereas vitamin E is a fat soluble antioxidant and secure our cell membrane from lipid peroxidation. The major sources of vitamin C in foods are grapefruits, oranges, lemon, tomato, sweet red peppers, cauliflower, sweet potato, black berry whereas sources of vitamin E are white beans, vegetable oils, sunflower seeds, almonds and peanuts. Carotenoids [23] defend us by quenching singlet oxygen oxy radicals and other ROS generated in our body. Best sources of carotenoids are carrots, cantaloupes, palm, apricots, mangoes, tomato and gac fruit (*Momordica*

cochinchinensis). Vegetables that deliver high content of polyphenol as antioxidants are spinach, cabbage, broccoli, beetroot, lettuce and asparagus. Most of the fruits like papaya, bananas, oranges, apple, and all varieties of melons, jackfruits, mango, grapes, guava, cherries and all varieties of berries including dried fruits such as raisins, figs and dates have notable contribution to supply polyphenol antioxidants in our body. Spices that have prominent role in our daily food-needs with adequate quantity of polyphenols as antioxidants are turmeric, garlic, clove, cinnamon, cumin, mustard seed, ginger, coriander, pepper, chilli powder, cardamom. Nuts such as cashew, walnuts, and pistachio also contribute modest polyphenol antioxidants.

Phytochemicals as antioxidants

Healthy food habits always keep the diseases at bay. However, it is quite common that it could not be accomplished routinely. Simultaneously, oxidative stress in course of time results in the deficiency of vital nutrients, minerals and antioxidants and as a whole our body instigates free radicals and ROS generation. It thus necessitates presence of adequate antioxidants to rescue our body from free radical damage. Keeping healthy and protecting our health from further deterioration, dietary antioxidant supplementations [24] are becoming a high priority. Phytochemicals/phytonutrients are found to be safe since ancient time possessing minimum side effects.

While it is impractical to list out all the redox complications relating various diseases in this short review, the authors are content to point out that at least in the cancerous situations several enzymes implicated in oxidative stress have abnormal expression. MsrA associated with reduction of S-stereoisomer of sulfoxide in methionine residues in enzymes is under expressed in human breast cancer [25] whereas over expression Heme Oxygenase (HO) potentiates cancerous growth in pancreatic cancer [26]. Similarly anti-oxidants can induce expression of detoxification phase-II enzymes in both neurons and astrocytes in neurodegenerative diseases. Importance of oxidative hypothesis in atherosclerosis has been well-known. Over oxidation in lipids leading to reactive carbonyls such as 4-hydroxynonenal (4-HNE), acrolein, malondialdehyde etc that ultimately react with DNA bases causing adverse effects; oxidation of essential proteins especially in their cysteine residues, DNA oxidation, down-regulation of Phase-II anti-oxidant enzymes, all-together contribute to disease generation and progression. Intervention in any of these processes by plant-derived anti-oxidants known for their safety and efficacy would be a welcome factor in mitigating these conditions. Some representative and promising candidates with high potential as antioxidant supplements are addressed as under.

Curcumin [27-29]: The major constituents of natural turmeric, namely curcuminoids, are known for their use as anti-inflammatory agent over centuries. Recent study reveals that intake of *curcumin* upto 12 g/day is safe with minimal toxicity [28]. Strong antioxidant effect of *curcumin*, a major property of curcuminoids, is responsible for not only their role as excellent anti-cancer agent but also for the prevention of other human diseases. The importance of the phenolic groups present in curcuminoids for the anti-oxidant activity is readily gleaned from the greatly diminished anti-oxidant activity of its mono- and di-glucuronide metabolites derived from *curcumin* [29].

Tetrahydrocurcumin [30,31]: This metabolite of *curcumin* has been found to be an effective antioxidant, surpassing *curcumin* in that respect and has been termed as a cascade anti-oxidant in the same way Carnosic acid-Rosmanol cascade has been referred [31]. Its antioxidant nature is found to be highly beneficial for slowing down the aging process and finds extensive use as one of the active cosmetic ingredients.

Tea polyphenols [(-)-Epigallocatechin gallate (EGCG), (-)-Epigallocatechin (EGC), (-)-Epicatechin gallate (ECG) and (-)-Epicatechin (EC)] [30,32-36]: These are important tea antioxidants with outstanding capability to quench the free radicals and subsequently to reduce the oxidative stress [32] and prevent cancer and cardiovascular diseases [33]. A recent paper on the use of green tea polyphenols brings out the impact of their anti-fatigue properties emphasized the importance of their anti-oxidant activity [34]. Its antioxidant effect is found to be beneficial for maintaining healthy eye [35,36].

Resveratrol [37-41]: Sources are red grapes, blue berries, red wine, dark chocolate and peanut skin. High content of resveratrol is found in Itadori tea. Itadori tea is mainly consumed in Japan and China and in use as natural medication [38]. It's wonderful antioxidant activity benefiting us from several diseases including cancer risk is summarized in recent reviews [39,40]. Recently synthetic resveratrol (>99%) (150 mg/day) was found safe in human health and has been approved by European Commission to use it as food supplement in form of tablet or capsule [41].

Garcinol [42,43]: Isolated from *Garcinia indica*. Garcinol is natural polyphenol with substantial antioxidant activity with potential of cancer prevention and several health benefits recently reviewed [43].

Pterostilbene [44-46]: This vital anti-oxidant is present in *Pterocarpus marsupium* extracts and is an excellent activator of Nrf2 transcription a responsible factor for several anti-oxidant enzymes. Pterostilbene is a better inhibitor of colon carcinogenesis. Its potent anti-diabetic activity is again ascribed to its anti-oxidant activity [45]. Its ability to inhibit an oxidative enzyme such as COX-2 was shown in a clinical trial [46].

Triterpenoids: Oleanolic acid [47,48], Ursolic acid [47] and Betulinic acid [49]: Oleanolic and Ursolic acids are effective inducers of metallothionein, a small cysteine-rich protein that acts like the anti-oxidant enzyme glutathione.

Flavonoids: Sources are in wide variety of vascular plants and also in fruits, vegetables, tea, and cocoa. Flavonoids enhance the antioxidants capacity of endogenous antioxidants [50].

Alkaloids [51-54]: Naturally occurring alkaloids possess antioxidant activity [51] with promising impact on health disorder [52]. They are found efficacious to use as anticancer agents [53,54].

Amla [55]: Amla is one of the most outstanding herbal products known since time immemorial. Its ability to attenuate oxidative stress has been well documented even though the consistent presence of the well-known anti-oxidant vitamin C as major constituent in the herbal extracts has been disputed [56]. β -Glucogallin (1-O-galloyl- β -D-glucose) is found one of its active antioxidant constituents with impressive photoprotection proficiency [57].

Boswellia serrata extract: *Boswellia serrata* gum resin is known as Indian frankincense [58] and in use as medicine for the treatment of several diseases since ancient times. Major constituents in *Boswellia serrata* include β -boswellic acid, keto β -boswellic acid and acetyl keto- β -boswellic acid. *Boswellia serrata* extract inhibits IL-1 β expression and it is believed that uncontrolled expression of IL-1 β results in the depletion of SOD leading to reduced collagen synthesis. Boswellic acids, especially acetyl-11-keto boswellic acid, are also inhibitors of 5-lipoxygenase, an oxidative enzyme implicated in leukotriene formation [59].

Salacinol: It is a novel, structurally unique anti-diabetic molecule isolated from *Salacia reticulata* documented in traditional medicine. Recent antioxidant analysis study reveals that *Salacia reticulata* possesses superior antioxidant activity [60,61] compared to other species of this

family [60]. It is concluded that better DPPH radical scavenging activity of *Salacia reticulata* is due to the presence of higher content of salacinol, kotalanol and magneferin in *Salacia reticulata* [60].

Forskolin: This diterpene natural molecule isolated from *Coleus forskohlii* reduces oxidative stresses via its antioxidant ability [62], possesses anti-diabetic activity [63] and presently in use as an anti-obesity preparations [64].

Conclusion

Antioxidants are mostly consumed through our diets; however the quantity may not be sufficient enough for our daily needs. This behooves to boost ourselves with supplementary anti-oxidants. Numerous studies have been performed demonstrating the potential benefit of antioxidants supplementation [24,59,65-67] despite concerns expressed on the use of excess antioxidants supplements [68].

References

- Dreher D, Junod AF (1996) Role of oxygen free radicals in cancer development. *Eur J Cancer* 32A: 30-38.
- Hussain SP, Hofseth LJ, Harris CC (2003) Radical causes of cancer. *Nat Rev Cancer* 3: 276-285.
- Singh R, Devi S, Rakesh G (2015) Role of free radical in atherosclerosis, diabetes and dyslipidaemia: larger-than-life. *Diabetes Metab Res Rev* 31:113-126.
- Watson JD (2014) Type 2 diabetes as a redox disease. *Lancet* 383: 841-843.
- Hadjigogos K (2003) The role of free radicals in the pathogenesis of rheumatoid arthritis. *Panminerva Med* 45: 7-13.
- Demir S, Yilmaz M, Koseoglu M, Akalin N, Aslan D, et al. (2003) Role of free radicals in peptic ulcer and gastritis. *Turk J Gastroenterol* 14: 39-43.
- Flamm ES, Demopoulos HB, Seligman ML, Poser RG, Ransohoff J (1978) Free radicals in cerebral ischemia. *Stroke* 9: 445-447.
- Maxwell SR, Lip GY (1997) Free radicals and antioxidants in cardiovascular disease. *Br J Clin Pharmacol* 44: 307-317.
- Kumar H, Lim HW, More SV, Kim BW, Koppula S, et al. (2012) The role of free radicals in the aging brain and Parkinson's Disease: convergence and parallelism. *Int J Mol Sci* 13:10478-10504.
- Fletcher AE (2010) Free radicals, antioxidants and eye diseases: evidence from epidemiological studies on cataract and age-related macular degeneration. *Ophthalmic Res* 44: 191-198.
- Thiagarajan R, Manikandan R (2013) Antioxidants and cataract. *Free Radic Res* 47: 337-345.
- Lynch ED, Kil J (2005) Compounds for the prevention and treatment of noise-induced hearing loss. *Drug Discov Today* 10: 1291-1298.
- Lynch ED, Gu R, Pierce C, Kil J (2004) Ebselen-mediated protection from single and repeated noise exposure in rat. *Laryngoscope* 114: 333-337.
- Nagy IZ (2001) On the true role of oxygen free radicals in the living state, aging, and degenerative disorders. *Ann NY Acad Sci* 928: 187-199.
- Halliwell B, Gutteridge JM (1995) The definition and measurement of antioxidants in biological systems. *Free Radic Biol Med* 18:125-126.
- Pisoschi AM, Pop A (2015) The role of antioxidants in the chemistry of oxidative stress: A review. *Eur J Med Chem* 97: 55-74.
- Roche M, Rondeau P, Singh NR, Tamus E, Bourdon E (2008) The antioxidant properties of serum albumin. *FEBS Letts* 582: 1783-1787.
- Apak R, Özyürek M, Güçlü K, Çapanoğlu E (2016) Antioxidant Activity/Capacity Measurement. 1. Classification, Physicochemical Principles, Mechanisms, and Electron Transfer (ET)-Based Assays. *J Agric Food Chem* 64: 997-1027.
- Apak R, Özyürek M, Güçlü K, Çapanoğlu E (2016) Antioxidant Activity/Capacity Measurement. 2. Hydrogen Atom Transfer (HAT)-Based, Mixed-Mode (Electron Transfer (ET)/HAT), and Lipid Peroxidation Assays. *J Agric Food Chem* 64: 1028-1045.
- Apak R, Özyürek M, Güçlü K, Çapanoğlu E (2016) Antioxidant Activity/Capacity Measurement. 3. Reactive Oxygen and Nitrogen Species (ROS/RNS) Scavenging Assays, Oxidative Stress Biomarkers, and Chromatographic/Chemometric Assays. *J Agric Food Chem* 64: 1046-1070.
- Carlsen MH, Halvorsen BL, Holte K, Bohn SK, Dragland S, et al. (2010) The total antioxidant content of more than 3100 foods, beverages, spices, herbs and supplements used worldwide. *Nutr J* 9: 3.
- Urquiza-Martínez MV, Navarro BF (2016) Antioxidant Capacity of Food. *Free Rad Antiox* 6: 1-12.
- Fiedor J, Burda K (2014) Potential Role of Carotenoids as Antioxidants in Human Health and Disease. *Nutrients* 6: 466-488.
- Fusco D, Colloca G, Lo Monaco MR, Cesari M (2007) Effects of antioxidant supplementation on the aging process. *Clin Interv Aging* 2: 377-387.
- De Luca A, Sanna F, Sallese M, Ruggiero C, Grossi M, et al. (2010) Methionine sulfoxide reductase A downregulation in human breast cancer cells results in a more aggressive phenotype. *Proc Natl Acad Sci USA* 107: 18628-18633.
- Sunamura M, Duda DG, Ghattas, MH, Lozonschi L, Motoi F, et al. (2003) Heme oxygenase-1 accelerates tumor angiogenesis of human pancreatic cancer. *Angiogenesis* 6: 15-24.
- Iqbal M, Sharma SD, Okazaki Y, Fujisawa M, Okada S (2003) Dietary supplementation of curcumin enhances antioxidant and phase II metabolizing enzymes in ddY male mice: possible role in protection against chemical carcinogenesis and toxicity. *Pharmacol Toxicol* 92: 33-38.
- Lao CD, Ruffin MT IV, Normolle D, Heath DD, Murray SI, et al. (2006) Dose escalation of a curcuminoid formulation. *BMC Complement Altern Med* 6: 10.
- Choudhury AK, Raja S, Mahapatra S, Nagabhushanam K, Majeed M (2015) Synthesis and Evaluation of the Anti-Oxidant Capacity of Curcumin Glucuronides, the Major Curcumin Metabolites. *Antioxidants* 4: 750-767.
- Kitani K, Osawa T, Yokozawa T (2007) The effects of tetrahydrocurcumin and green tea polyphenol on the survival of male C57BL/6 mice. *Biogerontology* 8: 567-573.
- Marrero JG, Andrés LS, Luis JG (2002) Semisynthesis of rosmanol and its derivatives. Easy access to abietatriene diterpenes isolated from the genus *Salvia* with biological activities. *J Nat Prod* 65: 986-989.
- Frei B, Higdon JV (2003) Antioxidant activity of tea polyphenols *in vivo*: evidence from animal studies. *J Nutr* 133: S3275-S3284.
- Khan N, Mukhtar H (2007) Tea polyphenols for health promotion. *Life Sci* 81: 519-533.
- Liudong F, Feng Z, Daoxing S, Xiufang Q, Xiaolong F, et al. (2011) Evaluation of anti-oxidant properties and anti-fatigue effect of green tea polyphenols. *Scientific Research and Essays* 6: 2624-2629.
- Rhone M, Basu A (2008) Phytochemicals and age-related eye diseases. *Nutr Rev* 66: 465-472.
- Zhang B, Safa R, Rusciano D, Osborne NN (2007) Epigallocatechin gallate, an active ingredient from green tea, attenuates damaging influences to the retina caused by ischemia/reperfusion. *Brain Res* 1159: 40-53.

37. Sahin K, Akdemir F, Orhan C, Tuzcu M, Hayirli A, et al. (2010) Effects of dietary resveratrol supplementation on egg production and antioxidant status. *Poult Sci* 89: 1190-1198.
38. Burns J, Yokota T, Ashihara H, Lean ME, Crozier A (2002) Plant foods and herbal sources of resveratrol. *J Agric Food Chem* 50: 3337-3340.
39. Yu W, Fu YC, Wang W (2012) Cellular and molecular effects of resveratrol in health and disease. *J Cell Biochem* 113: 752-759.
40. Baur JA, Sinclair DA (2006) Therapeutic potential of resveratrol: the *in vivo* evidence. *Nat Rev Drug Discov* 5: 493-506.
41. Scientific Opinion (2016) Safety of synthetic trans-resveratrol as a novel food pursuant to Regulation (EC) No 258/97. *EFSA Journal* 14: 4368.
42. Hung WL, Tsai ML, Sun PP, Tsai CY, Yang CC, et al. (2014) Protective effects of garcinol on dimethylnitrosamine-induced liver fibrosis in rats. *Food Funct* 5: 2883-2891.
43. Tang W, Pan MH, Sang S, Li S, Ho CT (2013) Garcinol from *Garcinia indica*: Chemistry and Health Beneficial effects. *ACS Symposium Series* 1129: 133-145.
44. Chiou YS, Tsai ML, Nagabhushanam K, Wang YJ, Wu CH, et al. (2011) Pterostilbene is more potent than resveratrol in preventing azoxymethane (AOM)-induced colon tumorigenesis via activation of the NF-E2-related factor 2 (Nrf2)-mediated antioxidant signaling pathway. *J Agric Food Chem* 59: 2725-2733.
45. Satheesh MA, Pari L (2006) The antioxidant role of pterostilbene in streptozotocin-nicotinamide-induced type 2 diabetes mellitus in Wistar rats. *J Pharm Pharmacol* 58: 1483-1490.
46. Hougee S, Faber J, Sanders A, de Jong RB, van den Berg WB, et al. (2005) Selective COX-2 inhibition by a *Pterocarpus marsupium* extract characterized by pterostilbene, and its activity in healthy human volunteers. *Planta Med* 71: 387-392.
47. Liu J (2005) Oleanolic and Ursolic acid: Research perspectives. *J Ethnopharmacol* 100: 92-94.
48. Wang X, Ye XL, Liu R, Chen HL, Bai H, et al. (2010) Antioxidant activities of Oleanolic acid *in vitro*: possible role of Nrf2 and MAP kinases. *Chem Biol Interact* 184: 328-337.
49. Yi J, Xia W, Wu J, Yuan L, Wu J, et al. (2014) Betulinic acid prevents alcohol-induced liver damage by improving the antioxidant system in mice. *J Vet Sci* 15: 141-148.
50. Pietta PG (2000) Flavonoids as antioxidants. *J Nat Prod* 63: 1035-1042.
51. Tiong SH, Looi CY, Hazni H, Arya A, Paydar M, et al. (2013) Antidiabetic and antioxidant properties of alkaloids from *Catharanthus roseus* (L.) G. Don. *Molecules* 18: 9770-9784.
52. Almagro L, Fernández-Pérez F, Pedreño MA (2015) Indole alkaloids from *Catharanthus roseus*: bioproduction and their effect on human health. *Molecules* 20: 2973-3000.
53. Mohan K, Jeyachandran R, Deepa (2012) Alkaloids as anticancer agents. *Annals of Phytomedicine* 1: 46-53.
54. Lu JJ, Bao JL, Chen XP, Huang M, Wang YT (2012) Alkaloids isolated from natural herbs as the anticancer agents. *Evid Based Complement Alternat Med* 2012: 485042.
55. Yang B, Liu P (2014) Composition and biological activities of hydrolyzable tannins of fruits of *Phyllanthus emblica*. *J Agric Food Chem* 62: 529-541.
56. Majeed M, Bhat B, Jadhav AN, Srivastava JS, Nagabhushanam K (2009) Ascorbic acid and tannins from *Emblca officinalis* Gaertn. Fruits—a revisit. *J Agric Food Chem* 57: 220-225.
57. Majeed M, Bhat B, Anand TSS (2010) Inhibition of UV induced adversaries by β -glucogallin from *Amla (Emblca officinalis)* Gaertn.) fruits. *Ind J Nat Prod Res* 1: 462-465.
58. Badria FA (2015) Frankincense (Heaven's Gift)—Chemistry, Biology, and Clinical Applications. In: Badria FA (Eds) Evidence-based Strategies in Herbal Medicine, Psychiatric Disorders and Emergency Medicine. *Intech* 1: 3-22.
59. Grover AK, Samson SE (2016) Benefits of antioxidant supplements for knee osteoarthritis: rationale and reality. *Nutr J* 15: 1.
60. Dias KLDK, Kottearachchi NS, Chandrasekara A, De Silva AN (2015) Antioxidant Analysis and In vitro Culture Establishment of Kothala Himbutu (*Salacia reticulata*). *Proceeding of 14th Agricultural Research Symposium* 1-5.
61. Yoshino K, Kanetaka T, Koga K (2015) Antioxidant Activity of *Salacia* Plant (*Salacia reticulata*). *Shokuhin Eiseigaku Zasshi* 56: 144-150.
62. Khatun S, Chatterjee NC, Cakilcioglu U (2011) Antioxidant activity of the medicinal plant *Coleus forskohlii* Briq. *Afr J Biotechnol* 10: 2530-2535.
63. Ríos-Silva M, Trujillo X, Trujillo-Hernández B, Sánchez-Pastor E, Urzúa Z, et al. (2014) Effect of chronic administration of forskolin on glycemia and oxidative stress in rats with and without experimental diabetes. *Int J Med Sci* 11: 448-452.
64. Majeed M (2012) *Coleus forskohlii* Extract in the Management of Obesity. In: Bagchi B, Preuss, HG (Eds) Obesity Epidemiology, Pathophysiology, and Prevention. 2nd Edition, CRC Press, Taylor & Francis Group, New York 703-728.
65. Myung SK, Kim Y, Ju W, Choi HJ, Choi Bae WK (2010) Effects of antioxidant supplements on cancer prevention: meta-analysis of randomized controlled trials. *Ann Oncol* 21: 166-179.
66. Myung SK, Ju W, Cho B, Oh SW, Park SM, et al. (2013) Efficacy of vitamin and antioxidant supplements in prevention of cardiovascular disease: systematic review and meta-analysis of randomised controlled trials. *BMJ* 346: f10.
67. Clifford T, Howatson G, West DJ, Stevenson EJ (2015) The Potential Benefits of Red Beetroot Supplementation in Health and Disease. *Nutrients* 7: 2801-2822.
68. Watson J (2013) Oxidants, antioxidants and the current incurability of metastatic cancers. *Open Biol* 3: 120144.