

# Immunological analysis of Malaria Infection in Carriers of Abnormal Hemoglobin EA or EE

Sourabié Y<sup>1,2\*</sup>, Bazié WW<sup>1</sup>, Sangaré I<sup>1,2</sup>, Sanou G<sup>5</sup>, Fumoux F<sup>4</sup> and Traoré Y<sup>3</sup>

<sup>1</sup>Centre Hospitalier Universitaire Sourô Sanou 01 BP 676, Bobo, 01 Burkina Faso

<sup>2</sup>Institut supérieur des sciences de la santé, Université polytechnique de Bobo Dioulasso, Burkina Faso

<sup>3</sup>Faculté des Sciences de Luminy, Aix Marseille II Université, France

<sup>4</sup>Unité de Formation et de Recherche des Sciences et Technologies, Université de Ouagadougou, Burkina Faso

<sup>5</sup>Centre National de Recherche et de Formation sur le Paludisme, Burkina Faso

\*Corresponding author: Yacouba Sourabié, Enseignant-chercheur en Immunologie à Institut supérieur des sciences de la santé, Université polytechnique de Bobo Dioulasso, Centre Hospitalo-universitaire Sourô Sanou, Service d'Immunologie et d'Hématologie, BP 676, Burkina Faso, Tel: 0022670710325; E-mail: [yacourabie@yahoo.fr](mailto:yacourabie@yahoo.fr)

Received date: 02 Nov 2015; Accepted date: 01 Apr 2016; Published date: 04 Apr 2016.

Citation: Sourabié Y, Bazié WW, Sangaré I, Sanou G, Fumoux F, et al. (2016) Immunological analysis of Malaria infection in carriers of abnormal Hemoglobin EA or EE. Int J Vaccine Immunizat 2(2): doi <http://dx.doi.org/10.16966/2470-9948.110>

Copyright: © 2016 Sourabié Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Background:** The purpose of our study is to establish the relationship between the rate of total immunoglobulin G in the sera of subjects with hemoglobin E or A genotype and resistance to infection by *Plasmodium falciparum*.

**Methods:** This is a prospective study over a period of five months on the sera of Vietnamese.

The serums were from whole blood collected in 2002 by venipuncture in subjects of study. The collected blood of each patient was centrifuged and the obtained serum was aliquoted into cryotubes and stored at 20°C for immunological studies (IgG assay). To maximize the sensitivity of ELISA test, we have coupled the antigenic peptides using glutaraldehyd with, highly polymerized poly-L-lysine as a resin. The enzymatic reaction then takes place in the dark for 30 min after addition of substrate before reading the results using a spectrophotometer (Opsys MR Dynex® Technology) set at the wave length of 405 nm.

**Results:** A total of 182 blood samples were taken from people belonging to 35 families. None of the subjects had clinical symptoms of malaria at the time of sampling. The distribution of the age is 3-79 years; the sex ratio (M/F) is 1.6. The proportions of hemoglobin are 42% of hemoglobin AA carriers and 58% of hemoglobin EA and EA carriers. In our different age groups, specific total IgG against antigenic peptides (MSP-1, MSP-2 and SR-11.1) of *P. falciparum* show an age-dependent distribution. In subjects 3-19 years, the titer of specific total IgG against *Plasmodium falciparum* is lower than that of more than 20 years. Comparison of specific IgG against the antigenic peptides MSP-1, MSP-2 and SR-11.1 of *P. falciparum* in the age groups, gives a significantly higher average for MSP-1 compared to SR-11.1 and MSP-2 (K<sub>hi</sub><sup>2</sup>=4.12) The rate of total IgG against the antigenic peptides at subjects with hemoglobin E carrier is higher than those of hemoglobin A (p=0.004).

**Conclusion:** We have shown that the response of specific total Ig G against the plasmodium antigens (MSP-1, MSP-2 and SR-11.1) is improved in patients over 20 years with hemoglobin E genotype compared to those with the AA genotype. In subjects under 20 years, no significant differences were obtained between the different groups of hemoglobin. MSP-2 and MSP-1 appear much more immunogenic than the SR11.1 in different age groups.

**Keywords:** Hemoglobin E; MSP1; MSP2; SR11.1; ELISA; IgG

## Introduction

In West Africa, it has been shown that after long exposure to intense transmission; it gradually develops a non-sterilizing immunity against *Plasmodium falciparum*. Moreover, in these endemic areas, some individuals appear naturally protected against clinical malaria and/or severe forms of malaria [1]. Suggesting a genetic control of resistance to malaria clinical infection, it has been clearly demonstrated in murine models that genetic factors control *Plasmodium* infection [2,3]. Studies have clearly demonstrated the role of abnormal hemoglobin in resistance to malaria attacks. In this resistance sickle cell trait (Hb AS) due to a change of the beta chain of glycin (6 beta: Glutamic acid → Valin) is the most studied genetic abnormalities of RBCs [4-6]. Thus, it was shown that susceptibility to *Plasmodium falciparum* infection and malaria attacks are less frequent in carriers of Hb S or C [7-9]. However, the mechanisms mediating this resistance remain unclear.

It was suggested that the protective effect in hemoglobinopathies may be partly due to the inhibition of *Plasmodium falciparum* increase in the

S parasitized erythrocytes [10,11] and might thus induce modulation of the immunoglobulin G response in carrier individuals of these hemoglobinopathies [4].

A recent proposal of protection against *Plasmodium falciparum* mechanism in the carrying hemoglobin C and S shows that it is linked to an antigen exposure PfEMP1 anomaly on the surface of infected sickle cell C or S [6]. Other immuno-epidemiological studies have also shown that the protection against malaria is directed at least partially by the antibody [12], with a high titer of Ig G directed against malarial antigens in situations Hb AS/SS/SC versus AA [9,13,14]. Thus, [15,16] showed that the cytophilic antibodies (IgG1 and IgG3) protect against *Plasmodium falciparum* by activating effectors cells through FcγRIIA while IgG4 and IgG2 block the protection mechanisms. And porting HbS, IgG3 response against MSP-2 may influence the decrease of the title of IgG2.

Another hemoglobin E (homozygous or heterozygous) due to a substitution of Glutamic acid for lysine at the 26th amino acid segment of A beta globin chain is frequent in South East Asia and has been

described as conferring resistance to multiplication in the erythrocytes of *Plasmodium falciparum* and *vivax* [10,17]. Chotivanich et al. [18] showed that patients had low parasite density in the carrying hemoglobin E which allowed protection against severe manifestations of malaria. Very few studies have been conducted to show the involvement of hemoglobin E in modulating the immune response against variant surface antigens of different stages of *Plasmodium falciparum* or *vivax*.

Data exist and have shown the role of hemoglobin E in the resistance of patients in severe malaria cases and simple. The purpose of our study is to establish the relationship between the title of total immunoglobulin G in the serum of subjects with hemoglobin E or A genotype and resistance to infection by *Plasmodium falciparum*.

## Methods

### Patients

This is a prospective study over a period of five months on the serum of Vietnamese living in the commune of XaTanh in Quang Tri's province and aged 3-79 years. The study took place at the Faculty of Pharmacy of the Timone, Marseille: Mixed Research Unit-MD3 (host-parasite relationship and immunogenetics Pharmacotherapeutic).

Our serums were from whole blood collected in 2002 by venipuncture in subjects of study. The collected blood of each patient was centrifuged and the obtained serum was aliquoted into cryotubes and stored at 20°C for immunological studies (IgG assay). The choice about the town Xa Thanh is justified by the fact that in this area there is a high prevalence of malaria and haemoglobinopathy E is high in this area.

### Laboratory Assays: ELISA after peptide coupling with glutaraldehyde

Direct sensitization by synthetic peptides on the ELISA plate is generally inefficient and highly dependent on their size and charge. To maximize the sensitivity of our test, we have coupled the antigenic peptides using glutaraldehyde using, highly polymerized poly-L-lysine as a resin.

**Principle:** poly-L-lysine is coated on the plate that serves as an anchor by exposing the NH<sub>3</sub><sup>+</sup> group in its side chains with reactive functions CHO glutaraldehyd. The covalent bond formed after chemical reaction is then used to fix the coupling agent to the plate. A second reaction between the second aldehyde function of glutaraldehyd and a primary amino group (NH<sub>3</sub>+terminal or lateral) allows covalent attachment of plasmodial antigens (MSP-1, MSP-2 and SR-11.1) to the poly-L-lysine and thus better attachment to the ELISA plate. The amino function of glycine was then used to block the aldehyde functions. We used the following protocol:

Poly-L-Lysine is coated on the plate overnight at 4°C (40 µg/ml in carbonate buffer pH=9.6), 100 µl/well. The plate is washed three times with PBS 1X and then 100 µl/well of 1% glutaraldehyd diluted in PBS 1X buffer were incubated for 30 minutes at room temperature. The plate is then washed three times and the antigenic peptide solution (10 µg/ml diluted in PBS 1X buffer) was added to the wells (100 µl/well). The plate is incubated overnight at room temperature, washed three times and then the free reactive functional groups are blocked by 1 M glycine (in PBS 1X, 200 µl/well) for 1 h at room temperature. Finally, the plated was washed and saturated with PBS-3% Milk (2 h at room temperature, 250 µl/well). Then we followed the steps of:

Each stage after the saturation of the plate is preceded by three washes in PBS 0.05% Tween, 200 µl per well.

Samples containing specific antibodies to be tested are diluted 1/20 in PBS 1X, for the measurement of specific total IgG.

For each plate, standards were included by using the plasma pool of African subjects as the positive controls. Negative controls are established

with self plasmas unexposed Europeans to *Plasmodium falciparum* and a "white" in which the plasma is replaced with PBS 1X dilution buffer. The standards were made by dilution in a range of ½.

The diluted plasma (1:20) are deposited at 100 µl/well and incubated for 2 h at room temperature.

Then the plate is coated with the goat's anti-human immunoglobulin G, a secondary antibody coupled to alkaline phosphatase (diluted 1/3000 in PBS 1X according to the manufacturer's recommendations (Beckman Coulter®) at 100 µl / well and 'incubated for 2 h at room temperature, stirring.

The development is carried out by depositing the substrate of the enzyme alkaline phosphatase, p-nitrophenyl phosphate (pNPP), prepared extemporaneously at the rate of 200 µl per well. The enzymatic reaction then takes place in the dark for 30 min after addition of substrate before reading the results using a spectrophotometer (OpsysMRDyner® Technology) set at the wave length of 405 nm.

### Statistical analysis of data

The processing and analysis of data is done using the Excel 2007 software for tables and graphs.

At all absorbance values, it was subtracted from the absorbance of "white".

To compare the values obtained between the various processes, we have standardized the values of the positive pool. This correction was then applied to the optical density of each serum.

The sample titration is carried out using a range obtained from pool of sera of Africans living in endemic areas. The following dilutions of factors were used: 20; 50; 100; 200 and 400. The measured absorbance for the 1/400 dilution was associated with the value of 10 AU/ml. The equation of the trend line obtained from the range allowed us to calculate for each absorbance measured its equivalent in AU/ml (Figures 1-3).

The tests used for the comparison of means and proportions were the tests of chi-squared and Fisher, Kruskal Wallis. P values <0.05 were statistically significant.

## Results

### Characteristics of the study population and the study area

Blood samples were taken in the town of Xa Thanh, Huong Hoa District. This town is located in a region above sea level, along the Sêpon river forms the border between Laos and Vietnam. The commune's territory covers 1200 ha and includes 9 hamlets. The population at the last census






Figure 1: The equation of the trend line obtained from the range with MSP-2



**Figure 2:** The equation of the trend line obtained from the range with SR-11.1



**Figure 3:** The equation of the trend line obtained from the range with MSP-1

has 2203 inhabitants and 382 families. All the inhabitants belong to ethnic Bru-Vân Kiêu. They are mainly farmers and herders. The families have on average 4.8 children, educational attainment remains low [19]. The Bru Van Kiêu practice intermittent crops, annually using burning zones remote from the permanent housing. They often spend the night in the forests. The habitat is mostly traditional; with the kitchen in the middle of the house (net use is limited due to fire fears). The people have very close relations with Laos and make frequent visits to Laos.

Huong Hoa district is area highly endemic malaria. More than 30 species of Anopheles (a total of 31 species in the province) have been described. The two main vectors of transmission are *An. Minimus* and *An. Dirus*. The majority of the catches of the province were carried out in the district of Huong Hoa (88% of *An. Minimus* and 95% of *An. Dirus*). The number of malaria cases is particularly strong during the rainy season (June to September). The data collected at the dispensary of Xa Thanh commune from 1997 to 2001 indicate an average of 125 cases of malaria per year for a population of 2203 people, with annual changes ranging from 72 to 194. Sixty percent of malaria cases are observed for four months, from June to September. There are *P. falciparum* infections (98%) and by *P. vivax* (2%). A total of 182 blood samples were taken from people belonging to 35 families. None of the subjects had clinical symptoms of malaria at the time of sampling. The distribution of the age is 3-79 years; the sex ratio (M/F) is 1.6.

### Distribution of subjects according type of hemoglobin and age

We left people in four age groups: children 0-10 years, teens 11-19 years, young adults aged 20-31 years and adults walls more than 32 years. For each age group, we have a similar proportion of individual AA and AE. The phenotypes of hemoglobin of these subjects were determined by electrophoresis. The proportions of hemoglobin are 42% of hemoglobin AA carriers and 58% of hemoglobin EA and EA carriers.

### Total immunoglobulin G rate according age

In our different age groups, specific total IgG against antigenic peptides (MSP-1, MSP-2 and SR-11.1) of *P. falciparum* show an age-dependent distribution. In subjects 3-19 years, the titer of specific total IgG against *Plasmodium falciparum* is lower than that of more than 20 years. Comparison of specific IgG against the antigenic peptides MSP-1, MSP-2 and SR-11.1 of *P. falciparum* in the age groups, gives a significantly higher average for MSP-1 compared to SR-11.1 and MSP-2 ( $\chi^2=4.12$ ) (Figure 4).

### Impact of hemoglobin on the rate of specific total IgG in patients from 3 to 19 years


The under 20 years are composed of 106 individuals whose average age is 11.7 years. The distribution of hemoglobin is 41/106 (39%) of hemoglobin AA carriers, (47/106) 44.3% of hemoglobin AE carriers and (16/106) 15.1% of hemoglobin EE carriers. In malaria's endemic areas, acquired immunity is required in subjects in their contacts with the antigen and is dependent on the level of transmission. We observed that the rate of specific total IgG against *P. falciparum* antigen such as MSP-1; MSP-2 and SR-11.1 obtained in the various groups of hemoglobin are not significantly different ( $p=0.07$ ) (Figures 4-6).

### Impact of hemoglobin on the specific IgG in subjects over 20 years


The rate of specific total IgG obtained in this age group are significantly different between groups of hemoglobin. Thus the rate of total IgG against the antigenic peptides at subjects with hemoglobin E carrier is higher than those of hemoglobin A (Figures 4-6). This difference was significant ( $p=0.004$ ). There is therefore a different acquisition of specific total IgG in subjects older than 20 years with hemoglobin E carrier.

### Discussion


In our study, we have observed that the rate of hemoglobin E is (58%). These results are higher than those found by Fucharoen G et al. [20] in Thailand (50%) and Flatz, et al. [21] Laos (43%).



**Figure 4:** The effect of age and hemoglobin on the rate of anti-MSP-1 total IgG



**Figure 5:** The effect of age and hemoglobin on the rate of anti-MSP-2 total IgG



**Figure 6:** The effect of age and hemoglobin on the rate of anti-SR-11.1 total IgG

Our data nonetheless confirm the strong representation of hemoglobinopathies E in Asia and its coincidence with the current distribution of malaria. Hemoglobin S is found in sub-Saharan Africa, in the equatorial belt, the Middle East and parts of India. The Hemoglobin C is common in the northwest of Africa where its distribution coincides with that of hemoglobin S. They are absent in Asia.

We also observed that the rate of specific IgG against MSP-1 levels is significantly higher than SR-11.1 and MSP-2 in the age groups. We have noted that the recognition of specific antibodies to the three peptides MSP-1, MSP-2 and SR-11.1 for the individuals in our study is low.

These low titers of antibodies specific for these antigens make it difficult to evaluate a clear relationship in the different groups of hemoglobin. This result could be explained by the variability of response options to many antigens of parasite strains. This could also be explained by the fact that IgG2 (and IgG4) are considered non cytophilic antibodies capable of blocking the protective mechanisms by preventing the recognition of antigens by antibodies cytophilic [22,23].

Three antigenic peptides we had used, MSP-1 and MSP-2 appear much more immunogenic than SR-11.1.

In our study, no significant differences between the titers of specific total IgG in hemoglobin groups have been found in subjects less than 20 years. Ntoumi et al. [15] in 2005 in Gabonese children with hemoglobin

AS showed a difference in antibody titers. This could be explained by the complex pathways of invasion of red blood cells by *P. falciparum* which increase the titers of IgG cytophilic against MSP-1 and MSP-2.

Our results could be explained by the fact that in endemic areas, the acquisition of antimalarial immunity required among young people depends on the level of transmission and is age-dependent [24,25]. In our population, subjects under 20 years have not yet reached the peak maximum amount of specific IgG.

We observed in the age group of over 20 years that the specific total IgG levels of antigenic peptides (MSP-1, MSP-2 and SR-11.1) are higher among carriers of hemoglobin E than hemoglobin AA carriers. This difference is significant. The same was done in Burkina Faso by Verra et al. [26], in 2007 which showed that the specific total IgG levels against the malarial antigens (MSP-1, MSP-2, MSP-3, EBA-175, and AMA-1) were higher when the subject was hemoglobin S or C carrier.

These results suggest that hemoglobin E carriers have better production of specific total IgG MSP-1; MSP-2 and SR-11 epitopes compared to normal hemoglobin AA carriers. It is possible that the multiplicity of *P. falciparum* genotypes contributes to increasing the recognition repertoire of *P. falciparum* and thus to accelerate the acquisition of protective immunity against the parasite.

## Conclusion

The analysis of our results shows that the frequency of hemoglobinopathies E (58%) higher in Vietnam. As we have shown that the response of specific total IgG against the plasmodium antigens (MSP-1, MSP-2 and SR-11.1) is improved in patients over 20 years with hemoglobin E genotype compared to those with the AA genotype. In subjects under 20 years, no significant differences were obtained between the different groups of hemoglobin. Even if we got a low rate of IgG against the antigenic peptides MSP-2 and MSP-1 appear much more immunogenic than the SR11.1 in different age groups.

## Author Contributions

Conceived and designed the experiments: SY TY FF. Performed the experiments: SY TY FF. Analyzed the data: SY TY,FF. Contributed reagents/materials/analysis tools: SY TY FF,SI BW. Wrote the paper: SY FF TY SI SG.

## Conflict of Interest

We declare that we have no conflict of interest.

## References

- Riley EM, Wagner GE, Ofori MF, Wheeler JG, Akanmori BD, et al. (2000) Lack of association between maternal antibody and protection of African infants from malaria infection. *Infect Immun* 68: 5856-5863.
- Ruwende C, Khoo SC, Snow RW, Yates SN, Kwiatkowski D, et al. (1995) Natural selection of hemi- and heterozygotes for G6PD deficiency in Africa by resistance to severe malaria. *Nature* 376: 246-249.
- McGuire W, Hill AV, Allsopp CE, Greenwood BM, Kwiatkowski D (1994) Variation in the TNF-alpha promoter region associated with susceptibility to cerebral malaria. *Nature* 371: 508-510.
- Marsh K, Otoo L, Hayes RJ, Carson DC, Greenwood BM (1989) Antibodies to blood stage antigens of *Plasmodium falciparum* in rural Gambians and their relation to protection against infection. *Trans R Soc Trop Med Hyg* 83: 293-303.
- Aidoo M, Terlouw DJ, Kolczak MS, McElroy PD, ter Kuile FO, et al. (2002) Protective effects of the sickle cell gene against malaria morbidity and mortality. *Lancet* 359: 1311-1312.

6. Cholera R, Brittain NJ, Gillrie MR, Lopera-Mesa TM, Diakité SA, et al. (2008) Impaired cytoadherence of *Plasmodium falciparum*-infected erythrocytes containing sickle hemoglobin. *Proc Natl Acad Sci* 105: 991-996.
7. Modiano D, Petrarca V, Sirima BS, Nebie I, Diallo D, et al. (1996) Different response to *Plasmodium falciparum* malaria in West African sympatric ethnic groups. *Proc Natl Acad Sci* 93: 13206-13211.
8. Modiano D, Luoni G, Sirima BS, Simporé J, Verra F, et al. (2001) Haemoglobin C protects against clinical *Plasmodium falciparum* malaria. *Nature* 414: 305-308.
9. Cabrera G, Cot M, Migot-Nabias F, Kremsner PG, Deloron P, et al. (2005) The sickle cell trait is associated with enhanced immunoglobulin G antibody responses to *Plasmodium falciparum* variant surface antigens. *J Infect Dis* 191: 1631-1638.
10. Nagel RL, Raventos-Suarez C, Fabry ME, Tanowitz H, Sicard D, et al. (1981) Impairment of the growth of *Plasmodium falciparum* in Hb EE erythrocytes. *J Clin Invest* 68: 303-305.
11. Chimma P, Roussillon C, Sratongno P, Ruangveerayuth R, Pattanapanyasat K, et al. (2009) A distinct peripheral blood monocyte phenotype is associated with parasite inhibitory activity in acute uncomplicated *Plasmodium falciparum* Malaria. *PLoS Pathog* 5: 631-1371.
12. Braga EM, Barros RM, Reis TA, Fontes CJ, Morais CG, et al. (2002) Association of the IgG response to *Plasmodium falciparum* Merozoite Surface Protein (C-terminal19 KD) with clinical immunity to malaria in to Brazilian amazon region. *Am J Trop Med Hyg* 5: 461-466.
13. Marsh K, Kinyanjui S (2006) Immune effectors mechanisms in malaria. *Parasite Immunol* 28: 51-60.
14. Ashley-Koch, Yang Q, Olney RS (2000) Sickle hemoglobin (HbS) allele and sickle cell disease: A HuGE Review. *Am J Epidemiol* 151: 839-845.
15. Ntoumi F, Flori L, Mayengue PI, Matondo Maya DW, Issifou S, et al. (2005) Influence of carriage of hemoglobin AS and the Fcγ Receptor IIa-R131 allele on levels of immunoglobulin G2 antibodies to *Plasmodium falciparum* Merozoite antigens in Gabonese children. *J Infect Dis* 192: 1975-1980.
16. Touré FS, Deloron P, Migot-Nabias F (2006) Analysis of human antibodies to erythrocyte binding antigen 175 peptide 4 of *Plasmodium falciparum*. *Clin Med Res* 4: 1-6.
17. Ohashi J, Naka I, Patarapotikul J, Hananantachai H, Brittenham G, et al. (2004) Extended linkage disequilibrium surrounding the hemoglobin E variant due to malarial selection. *Am J Hum Genet* 74: 1198-1208.
18. Chotivanich K, Udomsangpetch R, Pattanapanyasat K, Chierakul W, Simpson J, et al. (2002) Hemoglobin E: a balanced polymorphism protective against high parasitemia and thus severe P falciparum malaria. *Blood* 100: 1172-1176.
19. Archives de Documents (2001) Special Report: Mission FAO/WFP Crop and Evaluation of Food Supply in the Democratic Republic Lao People.
20. Fucharoen G, Fucharoen S, Sanchaisuriya K, Sae-ung N, Suyasunanond U, et al. (2002) Frequency distribution and haplotypic heterogeneity of βE-globin gene among eight minority groups of Northeast Thailand. *Hum Hered* 53: 18-22.
21. Flatz G, Sanguansermsri T, Sengchanh S, Horst D, Horst J (2004) The 'hot-spot' of Hb E [beta26 (B8) Glu-->Lys] in Southeast Asia: beta-globin anomalies in the Lao Theung population of southern Laos. *Hemoglobin* 28: 197-204.
22. Aucan C, Traoré Y, Tall F, Nacro B, Traoré-Leroux T, et al. (2000) High immunoglobulin G2 (IgG2) and low IgG4 levels are associated with human resistance to *Plasmodium falciparum* malaria. *Infect Immun* 68: 1252-1258.
23. Weatherall DJ, Miller LH, Baruch DI, Marsh K, Doumbo OK, et al. (2002) Malaria and the red cell. *Nature* 415: 673-679.
24. Williams TN, Mwangi TW, Roberts DJ, Alexander ND, Weatherall DJ, et al. (2005) An immune basis for malaria protection by the sickle cell trait. *PLoS Med* 2: e128.
25. Williams TN, Mwangi TW, Wambua S, Alexander ND, Kortok M, et al. (2005) Sickle cell trait and the risk of *Plasmodium falciparum* malaria and other childhood diseases. *J Infect Dis* 192: 178-186.
26. Verra F, Bancone G, Avellino P, Blot I, Simporé J, et al. (2007) Haemoglobin C and S role in acquired immunity against *Plasmodium falciparum* malaria. *PLoS One* 2: 978-1371.